



Gynecologic Cancer Intergroup (GCIG)

Cervix Cancer Committee

Thursday, May 30, 2013, 3:30 p.m. – 5:30 p.m.

LaSalle I Room, Doubletree Hotel, Chicago

CHAIR SATORU SAGAE.
CO-CHAIR: BRADLEY J. MONK
PAST-CHAIR WILLIAM SMALL, Jr.

*****Please sign Attendance Forms*****

WELCOME & INTRODUCTIONS (& COI declarations):

Satoru Sagae

RECENT CERVIX GCIG TRIALS THAT WERE CLOSED, PUBLISHED OR PRESENTED:

Publications:

- Lindsay R, Paul J, Siddiqui N, Davis J, Gaffney DK. Survey on the management of early cervical cancer among members of the GCIG. Int J Gynecol Cancer. 2012 Nov;22(9):1617-23.

Closed:

- GOG 240 (MANGO, NSGO): Cis/Paclitaxel +/- Bev vs Topo/Paclitaxel +/- Bev in stage IVB cervix cancer (Tewari) [Topo/Paclitaxel discontinued after planned fertility analysis] enrollment completed January 3, 2012 Monk

CURRENT ACTIVE/NEAR ACTIVATION RANDOMIZED TRIALS WITH GCIG PARTICIPATION:

- KGOG-Thai TACO: (RTOG, GICOM) (**CCRN** - Viet Nam, Thailand) Randomized Clinical Trial of Weekly versus Tri-Weekly Cisplatin based Chemoradiation in Locally Advanced Cervical Cancer S Y Ryu
- ANZGOG OUTBACK (GOG, RTOG) (**CCRN** - India) : Chemoradiotherapy +/- adjuvant chemotherapy Mileskin
- NCRI: INTERLACE: (**CCRN** – Romania, Belarus) Induction ChemoTherApy in Locally Advanced Cervical Cancer. McCormack
- NCIC-CTG CX 5 SHAPE: (DGOG,) (**CCRN --**) A RANDOMIZED TRIAL COMPARING RADICAL HYSTERECTOMY AND PELVIC NODE DISSECTION VS SIMPLE HYSTERECTOMY AND PELVIC NODE DISSECTION IN PATIENTS WITH LOW RISK EARLY STAGE CX CA: Fyles for Plante
- GOG-0278 Conservative surgery (cone/nodes or hyst/(nodes) in early stage cervical cancer. (evaluation of physical function and QOL) Monk for Covens
- GOG-0279 Phase II Trial Evaluating Cisplatin (NSC #119875) and Gemcitabine (NSC #613327) Concurrent with Intensity-Modulated Radiation Therapy (IMRT) in Treatment of Locally Adv. Squamous Cell Carcinoma of the Vulva Monk
- KGOG-0801/GOG 263 (RTOG): RT vs CRT in intermediate risk cervix cancer after hysterectomy SY Ryu
- ACRIN 6671(GOG, KGOG, JGOG) Utility of Preoperative FDG-PET/CT Prior to Primary Chemoradiation Therapy to Detect Retroperitoneal Lymph Node Metastasis in Patients With Locoregionally Advanced Carcinoma of the Cervix (IB2, IIA =4 CM, IIB-IVA) or Endometrium (Grade 3 Endometrioid Endometrial Carcinoma; Serous Papillary Carcinoma, Clear Cell Carcinoma, or Carcinosarcoma (Any Grade); and Grade 1 or 2 Endometrioid Endometrial Carcinoma With Cervical Stromal Involvement Overt in Clinical Examination or Confirmed by Endocervical Curettage) Monk for S. Lee
- RTOG-0724 (GOG): ChemoRT with and without adjuvant chemotherapy in high risk cervix cancer after hysterectomy Jhingran
- RTOG: A RANDOMIZED PHASE III STUDY OF STANDARD VS. IMRT PELVIC RADIATION FOR POST-OPERATIVE TREATMENT OF ENDOMETRIAL AND CERVICAL CANCER: TIME-C Small

- DGOG GROINSS-V II (EORTC, GOG) : Groningen International Study on Sentinel Nodes in Vulvar Cancer Creutzberg for van der Zee
- EORTC 55994: Randomized phase III study of neoadjuvant chemotherapy followed by surgery vs. concomitant radiotherapy and chemotherapy in FIGO stage Ib2, IIa>4 cm or IIb cervical cancer Casado

PROPOSED OR DEVELOPING CONCEPTS:

DISCUSSIONS

- Cervix Cancer Research Network (CCRN): (Gaffney/Kitchener)

ADJOURN

Of Note:

2:00pm CCRN satellite session ---- State II Room