


**A RANDOMIZED TRIAL COMPARING RADICAL HYSTERECTOMY AND PELVIC
NODE DISSECTION VS SIMPLE HYSTERECTOMY AND PELVIC NODE DISSECTION
IN PATIENTS WITH LOW RISK EARLY STAGE CERVICAL CANCER**

A Gynecologic Cancer Intergroup (GCIg) Trial led by the NCIC CTG

GCIg Trial Designation: The **SHAPE Trial
NCIC CTG Protocol Number: **CX.5****

Chair: Marie Plante

**NCIC CTG
NCIC GEC**



SHAPE-UPDATE

∞ Centers opened in **Canada: 13**

∞ Accrual to date: **11**

∞ **International groups status**

- **The Netherlands** have received **approval and funding !**
- **UK Group** has received approval for funding and will be opened to accrual by Spring 2014
- **Austria** has received approval and will activate by the end of 2013.
- **France and Germany** will activate by Spring 2014.
- **Ireland** will activate by the end of 2013.
- **The Nordic Society** will activate by the end of 2013.
- **The Belgium Group** will activate by the end of 2013.
- **Mexico, Korea, China and COGI (the group at Stanford Uni): ???**

SHAPE-UPDATE

∞ Adjuvant treatment

- Will be **left at the discretion** of each center based on local policy
- Specific references to **Cisplatin** administration has been removed from the protocol
- This amendment has been submitted to **Health Canada** (October 2013)

SHAPE-UPDATE

- ❧ Simplified randomization process
 - Maximum of **20 weeks** from diagnosis
- ❧ The **surgical approach** is no longer a stratification factor
- ❧ All patients must undergo **PLND**
 - SLN optional
 - LSG optional (preferred)